

ANALYSIS OF VOLATILE COMPOUNDS IN CORIANDER



**INSTITUT PENYELIDIKAN, PEMBANGUNAN & PENGKOMERSILAN
(IRDC)
UNIVERSITI TEKNOLOGI MARA
40450 SHAH ALAM
MALAYSIA**


PREPARED BY:

**ROZITA OSMAN
ASSOCIATE PROF. DR NOR'ASHIKIN SAIM
NORAINI HAMZAH**

FEBRUARI 2006

PROJECT TEAM MEMBERS

ROZITA OSMAN
Project Leader



.....
Tandatangan

ASSOCIATE PROF. DR NOR'ASHIKIN SAIM
Project Member



.....
Tandatangan

NORAINI HAMZAH
Project Member



.....
Tandatangan

TABLE OF CONTENTS

	Page
SURAT PERLANTIKAN	ii
SURAT PENYERAHAN LAPORAN	iii
PROJECT TEAM MEMBERS	iv
ACKNOWLEDGEMENT	v
TABLE OF CONTENTS	vi
LIST OF TABLES	viii
LIST OF FIGURES	ix
ABSTRACT	x
CHAPTER 1 INTRODUCTION	1
CHAPTER 2 LITERATURE REVIEW	3
2.1 Coriander (<i>Coriandrum sativum</i> L)	3
2.2 Solid Phase Microextraction (SPME)	5
2.3 Application of SPME in the Analysis of Volatile Compounds	7
2.4 Subcritical Water Extraction (SWE)	7
CHAPTER 3 EXPERIMENTAL	10
3.1 Reagents	10
3.2 Instrumentation	10
3.2.1 Solid Phase Microextraction (SPME)	10
3.2.2 Gas Chromatography-Mass Spectrometry (GC-MSD)	10
3.2.3 Accelerated Solvent Extraction (Dionex ASE 200)	10
3.3 Coriander Samples	11

3.4	SPME Analysis of Coriander	11
3.5	SPME Procedure	11
3.6	Subcritical Water Extraction (SWE)	11
3.7	Hydrodistillation	13
3.8	Analysis by Gas Chromatography-Mass Spectrometry Detector (GC-MSD)	13
CHAPTER 4	RESULTS AND DISCUSSION	15
4.1	SPME Performance	15
4.2	Optimization of SPME Operating Parameters	15
4.2.1	Effect of Extraction Temperature	15
4.2.2	Effect of Extraction Time	17
4.2.3	Effect of Desorption Time	18
4.2.4	Optimum SPME Extraction Conditions for Volatile Compounds in Coriander	20
4.3	Extraction and Identification of Volatiles from Coriander Leaves, Seeds and Powder using SPME	20
4.3.1	Volatiles from Leaves	20
4.3.2	Volatiles from Seeds	23
4.3.3	Volatiles from Powder	26
4.3.4	Comparison between the profile of Volatiles of Coriander Seeds and Coriander Powder	26
4.4	Extraction of Essential Oil by Subcritical Water Extraction (SWE)	27
4.4.1	Effect of Extraction Temperature	30
4.4.2	Effect of Extraction Pressure	31
4.4.3	Comparison of SWE and Hydrodistillation (HD)	31

ABSTRACT

This study focuses on the analysis of volatile compounds in coriander leaves and seeds using solid-phase microextraction (SPME) and subcritical water extraction (SWE). Separation and identification of the compounds were done using gas chromatography with mass spectrometry detector (GC-MSD). Optimization of SPME operating parameters (extraction time, temperature and desorption time) were studied using 85 μm polyacrylate fiber. Optimum SPME conditions of 40 minutes extraction time, 65 $^{\circ}\text{C}$ extraction temperature and desorption time of 130 seconds were used to analyze fresh coriander leaves, seeds and powder. It was found that the profile of volatile compounds from these samples differs, resulting in the different flavour. SPME combined with GC-MSD can provide a simple and fast method in profiling volatile compounds in coriander. In the application of SWE in extracting essential oil from coriander seeds, the effects of extraction time and extraction temperature on the quantity and quality of the essential oil were evaluated. The SWE method was compared with hydrodistillation method and it was found that the efficiency (g oil/g of coriander) of SWE was 5 times higher than that provided by hydrodistillation with reduced extraction time. In addition, using SWE the composition of the oil can be manipulated by selecting the temperature and adjusting the pressure.